
### Pre- and Post-Survey Results




## Ability to identify damage mechanisms included in the traditional understanding of MS pathophysiology


Note: Highlighted p-values are statistically significant




# Ability to identify that long-term disability, axonal damage, and response to therapy are positively affected by the early recognition and treatment of MS




## Ability to consider specific patient characteristics and identify actions and goals recommended by the CMSC that can help empower patients with MS




#### Ability to consider specific patient characteristics and identify the most appropriate first-line treatment for a patient with MS



## Ability to identify clinical indicators of suboptimal response to treatment




#### Ability to identify serious safety concerns that are associated with emerging MS therapies



#### Pre-, Post-, and 30-day Survey Results




### Ability to identify damage mechanisms included in the traditional understanding of MS pathophysiology




Pre to post: P = 0.002

Post to 30-day: P = 1.000




# Ability to identify that long-term disability, axonal damage, and response to therapy are positively affected by the early recognition and treatment of MS




Pre to post: P = 0.698

Post to 30-day: P = 0.188




## Ability to consider specific patient characteristics and identify the most appropriate first-line treatment for a patient with MS




Pre to post: *P* < 0.001

Post to 30-day: P = 0.383




### Ability to identify clinical indicators of suboptimal response to treatment




Pre to post: P = 0.012

Post to 30-day: P = 0.922




## Ability to identify serious safety concerns that are associated with emerging MS therapies



Pre to post: P = 0.020

Post to 30-day: P = 0.166



**Limited Sample**